

Certain Facts about Families of Subsets of Many Sorted Sets

Artur Korniłowicz
Institute of Mathematics
Warsaw University
Białystok

MML Identifier: MSSUBFAM.

WWW: <http://mizar.org/JFM/Vol7/mssubfam.html>

The articles [18], [5], [21], [16], [22], [3], [1], [4], [7], [6], [19], [20], [2], [12], [13], [14], [8], [15], [10], [17], [9], and [11] provide the notation and terminology for this paper.

1. PRELIMINARIES

For simplicity, we adopt the following rules: I, G, H, i denote sets, A, B, M denote many sorted sets indexed by I , s_1, s_2, s_3 denote families of subsets of I , v, w denote subsets of I , and F denotes a many sorted function indexed by I .

The scheme *MSFExFunc* deals with a set \mathcal{A} , many sorted sets \mathcal{B}, \mathcal{C} indexed by \mathcal{A} , and a ternary predicate \mathcal{P} , and states that:

There exists a many sorted function F from \mathcal{B} into \mathcal{C} such that for every set i if $i \in \mathcal{A}$, then there exists a function f from $\mathcal{B}(i)$ into $\mathcal{C}(i)$ such that $f = F(i)$ and for every set x such that $x \in \mathcal{B}(i)$ holds $\mathcal{P}[f(x), x, i]$

provided the following condition is met:

- For every set i such that $i \in \mathcal{A}$ and for every set x such that $x \in \mathcal{B}(i)$ there exists a set y such that $y \in \mathcal{C}(i)$ and $\mathcal{P}[y, x, i]$.

Next we state a number of propositions:

- (1) If $s_1 \neq \emptyset$, then $\text{Intersect}(s_1) \subseteq \bigcup s_1$.
- (2) If $G \in s_1$, then $\text{Intersect}(s_1) \subseteq G$.
- (3) If $\emptyset \in s_1$, then $\text{Intersect}(s_1) = \emptyset$.
- (4) For every subset Z of I such that for every set Z_1 such that $Z_1 \in s_1$ holds $Z \subseteq Z_1$ holds $Z \subseteq \text{Intersect}(s_1)$.
- (5) If $s_1 \neq \emptyset$ and for every set Z_1 such that $Z_1 \in s_1$ holds $G \subseteq Z_1$, then $G \subseteq \text{Intersect}(s_1)$.
- (6) If $G \in s_1$ and $G \subseteq H$, then $\text{Intersect}(s_1) \subseteq H$.
- (7) If $G \in s_1$ and G misses H , then $\text{Intersect}(s_1)$ misses H .
- (8) If $s_3 = s_1 \cup s_2$, then $\text{Intersect}(s_3) = \text{Intersect}(s_1) \cap \text{Intersect}(s_2)$.
- (9) If $s_1 = \{v\}$, then $\text{Intersect}(s_1) = v$.

- (10) If $s_1 = \{v, w\}$, then $\text{Intersect}(s_1) = v \cap w$.
- (11) If $A \in B$, then A is an element of B .
- (12) For every non-empty many sorted set B indexed by I such that A is an element of B holds $A \in B$.
- (13) For every function f such that $i \in I$ and $f = F(i)$ holds $(\text{rng}_\kappa F(\kappa))(i) = \text{rng } f$.
- (14) For every function f such that $i \in I$ and $f = F(i)$ holds $(\text{dom}_\kappa F(\kappa))(i) = \text{dom } f$.
- (15) For all many sorted functions F, G indexed by I holds $G \circ F$ is a many sorted function indexed by I .
- (16) Let A be a non-empty many sorted set indexed by I and F be a many sorted function from A into $\mathbf{0}_I$. Then $F = \mathbf{0}_I$.
- (17) If A is transformable to B and F is a many sorted function from A into B , then $\text{dom}_\kappa F(\kappa) = A$ and $\text{rng}_\kappa F(\kappa) \subseteq B$.

2. FINITE MANY SORTED SETS

Let us consider I . Observe that every many sorted set indexed by I which is empty yielding is also locally-finite.

Let us consider I . Observe that $\mathbf{0}_I$ is empty yielding and locally-finite.

Let us consider I, A . Note that there exists a many sorted subset indexed by A which is empty yielding and locally-finite.

One can prove the following proposition

- (18) If $A \subseteq B$ and B is locally-finite, then A is locally-finite.

Let us consider I and let A be a locally-finite many sorted set indexed by I . Note that every many sorted subset indexed by A is locally-finite.

Let us consider I and let A, B be locally-finite many sorted sets indexed by I . One can verify that $A \cup B$ is locally-finite.

Let us consider I, A and let B be a locally-finite many sorted set indexed by I . Note that $A \cap B$ is locally-finite.

Let us consider I, B and let A be a locally-finite many sorted set indexed by I . One can verify that $A \cap B$ is locally-finite.

Let us consider I, B and let A be a locally-finite many sorted set indexed by I . One can check that $A \setminus B$ is locally-finite.

Let us consider I, F and let A be a locally-finite many sorted set indexed by I . Observe that $F^\circ A$ is locally-finite.

Let us consider I and let A, B be locally-finite many sorted sets indexed by I . Observe that $\llbracket A, B \rrbracket$ is locally-finite.

Next we state three propositions:

- (19) If B is non-empty and $\llbracket A, B \rrbracket$ is locally-finite, then A is locally-finite.
- (20) If A is non-empty and $\llbracket A, B \rrbracket$ is locally-finite, then B is locally-finite.
- (21) A is locally-finite iff 2^A is locally-finite.

Let us consider I and let M be a locally-finite many sorted set indexed by I . Note that 2^M is locally-finite.

One can prove the following four propositions:

- (22) Let A be a non-empty many sorted set indexed by I . Suppose A is locally-finite and for every many sorted set M indexed by I such that $M \in A$ holds M is locally-finite. Then $\bigcup A$ is locally-finite.

- (23) If $\bigcup A$ is locally-finite, then A is locally-finite and for every M such that $M \in A$ holds M is locally-finite.
- (24) If $\text{dom}_\kappa F(\kappa)$ is locally-finite, then $\text{rng}_\kappa F(\kappa)$ is locally-finite.
- (25) Suppose $A \subseteq \text{rng}_\kappa F(\kappa)$ and for every set i and for every function f such that $i \in I$ and $f = F(i)$ holds $f^{-1}(A(i))$ is finite. Then A is locally-finite.

Let us consider I and let A, B be locally-finite many sorted sets indexed by I . Observe that $\text{MSFuncs}(A, B)$ is locally-finite.

Let us consider I and let A, B be locally-finite many sorted sets indexed by I . One can verify that $A \dot{-} B$ is locally-finite.

In the sequel X, Y, Z denote many sorted sets indexed by I .

The following propositions are true:

- (26) Suppose X is locally-finite and $X \subseteq \llbracket Y, Z \rrbracket$. Then there exist A, B such that A is locally-finite and $A \subseteq Y$ and B is locally-finite and $B \subseteq Z$ and $X \subseteq \llbracket A, B \rrbracket$.
- (27) Suppose X is locally-finite and Z is locally-finite and $X \subseteq \llbracket Y, Z \rrbracket$. Then there exists A such that A is locally-finite and $A \subseteq Y$ and $X \subseteq \llbracket A, Z \rrbracket$.
- (28) Let M be a non-empty locally-finite many sorted set indexed by I . Suppose that for all many sorted sets A, B indexed by I such that $A \in M$ and $B \in M$ holds $A \subseteq B$ or $B \subseteq A$. Then there exists a many sorted set m indexed by I such that $m \in M$ and for every many sorted set K indexed by I such that $K \in M$ holds $m \subseteq K$.
- (29) Let M be a non-empty locally-finite many sorted set indexed by I . Suppose that for all many sorted sets A, B indexed by I such that $A \in M$ and $B \in M$ holds $A \subseteq B$ or $B \subseteq A$. Then there exists a many sorted set m indexed by I such that $m \in M$ and for every many sorted set K indexed by I such that $K \in M$ holds $K \subseteq m$.
- (30) If Z is locally-finite and $Z \subseteq \text{rng}_\kappa F(\kappa)$, then there exists Y such that $Y \subseteq \text{dom}_\kappa F(\kappa)$ and Y is locally-finite and $F^\circ Y = Z$.

3. A FAMILY OF SUBSETS OF MANY SORTED SETS

Let us consider I, M . A subset family of M is a many sorted subset indexed by 2^M .

Let us consider I, M . Note that there exists a subset family of M which is non-empty.

Let us consider I, M . Then 2^M is a subset family of M .

Let us consider I, M . Observe that there exists a subset family of M which is empty yielding and locally-finite.

One can prove the following proposition

- (31) $\mathbf{0}_I$ is an empty yielding locally-finite subset family of M .

Let us consider I and let M be a locally-finite many sorted set indexed by I . Observe that there exists a subset family of M which is non-empty and locally-finite.

We adopt the following convention: S_1, S_2, S_3 are subset families of M , S_4 is a non-empty subset family of M , and V, W are many sorted subsets indexed by M .

Let I be a non empty set, let M be a many sorted set indexed by I , let S_1 be a subset family of M , and let i be an element of I . Then $S_1(i)$ is a family of subsets of $M(i)$.

One can prove the following propositions:

- (32) If $i \in I$, then $S_1(i)$ is a family of subsets of $M(i)$.
- (33) If $A \in S_1$, then A is a many sorted subset indexed by M .
- (34) $S_1 \cup S_2$ is a subset family of M .
- (35) $S_1 \cap S_2$ is a subset family of M .

- (36) $S_1 \setminus A$ is a subset family of M .
- (37) $S_1 \dot{-} S_2$ is a subset family of M .
- (38) If $A \subseteq M$, then $\{A\}$ is a subset family of M .
- (39) If $A \subseteq M$ and $B \subseteq M$, then $\{A, B\}$ is a subset family of M .
- (40) $\bigcup S_1 \subseteq M$.

4. INTERSECTION OF A FAMILY OF MANY SORTED SETS

Let us consider I, M, S_1 . The functor $\bigcap S_1$ yielding a many sorted set indexed by I is defined by:

(Def. 2)¹ For every set i such that $i \in I$ there exists a family Q of subsets of $M(i)$ such that $Q = S_1(i)$ and $(\bigcap S_1)(i) = \text{Intersect}(Q)$.

Let us consider I, M, S_1 . Then $\bigcap S_1$ is a many sorted subset indexed by M .

The following propositions are true:

- (41) If $S_1 = \mathbf{0}_I$, then $\bigcap S_1 = M$.
- (42) $\bigcap S_4 \subseteq \bigcup S_4$.
- (43) If $A \in S_1$, then $\bigcap S_1 \subseteq A$.
- (44) If $\mathbf{0}_I \in S_1$, then $\bigcap S_1 = \mathbf{0}_I$.
- (45) Let Z, M be many sorted sets indexed by I and S_1 be a non-empty subset family of M . Suppose that for every many sorted set Z_1 indexed by I such that $Z_1 \in S_1$ holds $Z \subseteq Z_1$. Then $Z \subseteq \bigcap S_1$.
- (46) If $S_1 \subseteq S_2$, then $\bigcap S_2 \subseteq \bigcap S_1$.
- (47) If $A \in S_1$ and $A \subseteq B$, then $\bigcap S_1 \subseteq B$.
- (48) If $A \in S_1$ and $A \cap B = \mathbf{0}_I$, then $\bigcap S_1 \cap B = \mathbf{0}_I$.
- (49) If $S_3 = S_1 \cup S_2$, then $\bigcap S_3 = \bigcap S_1 \cap \bigcap S_2$.
- (50) If $S_1 = \{V\}$, then $\bigcap S_1 = V$.
- (51) If $S_1 = \{V, W\}$, then $\bigcap S_1 = V \cap W$.
- (52) If $A \in \bigcap S_1$, then for every B such that $B \in S_1$ holds $A \in B$.
- (53) Let A, M be many sorted sets indexed by I and S_1 be a non-empty subset family of M . Suppose $A \in M$ and for every many sorted set B indexed by I such that $B \in S_1$ holds $A \in B$. Then $A \in \bigcap S_1$.

Let us consider I, M and let I_1 be a subset family of M . We say that I_1 is additive if and only if:

(Def. 3) For all A, B such that $A \in I_1$ and $B \in I_1$ holds $A \cup B \in I_1$.

We say that I_1 is absolutely-additive if and only if:

(Def. 4) For every subset family F of M such that $F \subseteq I_1$ holds $\bigcup F \in I_1$.

We say that I_1 is multiplicative if and only if:

(Def. 5) For all A, B such that $A \in I_1$ and $B \in I_1$ holds $A \cap B \in I_1$.

We say that I_1 is absolutely-multiplicative if and only if:

¹ The definition (Def. 1) has been removed.

(Def. 6) For every subset family F of M such that $F \subseteq I_1$ holds $\bigcap F \in I_1$.

We say that I_1 is properly upper bound if and only if:

(Def. 7) $M \in I_1$.

We say that I_1 is properly lower bound if and only if:

(Def. 8) $\mathbf{0}_I \in I_1$.

Let us consider I, M . Observe that there exists a subset family of M which is non-empty, additive, absolutely-additive, multiplicative, absolutely-multiplicative, properly upper bound, and properly lower bound.

Let us consider I, M . Then 2^M is an additive absolutely-additive multiplicative absolutely-multiplicative properly upper bound properly lower bound subset family of M .

Let us consider I, M . Note that every subset family of M which is absolutely-additive is also additive.

Let us consider I, M . Observe that every subset family of M which is absolutely-multiplicative is also multiplicative.

Let us consider I, M . Observe that every subset family of M which is absolutely-multiplicative is also properly upper bound.

Let us consider I, M . Note that every subset family of M which is properly upper bound is also non-empty.

Let us consider I, M . Note that every subset family of M which is absolutely-additive is also properly lower bound.

Let us consider I, M . One can verify that every subset family of M which is properly lower bound is also non-empty.

REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinall.html>.
- [2] Ewa Burakowska. Subalgebras of many sorted algebra. Lattice of subalgebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_2.html.
- [3] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [4] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [5] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [6] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_4.html.
- [7] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [8] Artur Kornilowicz. On the group of automorphisms of universal algebra and many sorted algebra. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/autalg_1.html.
- [9] Artur Kornilowicz. Definitions and basic properties of boolean and union of many sorted sets. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/mboolean.html>.
- [10] Artur Kornilowicz. Extensions of mappings on generator set. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/extens_1.html.
- [11] Artur Kornilowicz. Some basic properties of many sorted sets. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/pzfmisc1.html>.
- [12] Małgorzata Korolkiewicz. Homomorphisms of many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_3.html.
- [13] Beata Madras. Product of family of universal algebras. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/pralg_1.html.
- [14] Beata Madras. Products of many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/pralg_2.html.

- [15] Yatsuka Nakamura, Piotr Rudnicki, Andrzej Trybulec, and Pauline N. Kawamoto. Preliminaries to circuits, I. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/pre_circ.html.
- [16] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/setfam_1.html.
- [17] Alexander Yu. Shibakov and Andrzej Trybulec. The Cantor set. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/cantor_1.html.
- [18] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [19] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pboole.html>.
- [20] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_1.html.
- [21] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [22] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received October 27, 1995

Published January 2, 2004
